

Rotating Squares Look Like Pincushions

i-Perception

September-October 2016, 1–4

© The Author(s) 2016

DOI: 10.1177/2041669516664741

ipe.sagepub.com



Stuart Anstis

Department of Psychology, University of California, San Diego, USA

Sae Kaneko

Research Institute of Electrical Communication, Tohoku University, Japan;
Department of Psychology, University of California, San Diego, USA

Abstract

Rotating squares appeared to be distorted into pincushions with concave sides. These illusory shape changes were caused by a perceived compression along the curved path of motion.

Keywords

motion perception, illusion, shapes or objects

The outline of a square that rotates about its own center at 0.5 to 1 rev/s, appears to be distorted into a pincushion with pointy corners and concave sides. This is not simply a display artefact, since a square printed on paper shows pincushioning when spun on a turntable.

We can define the distortion ratio of a pincushion as a percentage decrement—the ratio of its pinched width to the width of an undistorted square (Figure 1).

We noticed that nested concentric squares give a strong effect as compared with a single square (online Movie 1). The corners appear to stick out too far, and the straight sides of the square look curved inwards. We also noticed that the amount of apparent pincushion distortion increases with rotation rate (up to about 1.5 rev/s, when computer aliasing breaks up the pattern).

We measured the pincushion distortion by a nulling method. We noticed that the extent of “pincushioning” increased with angular velocity. So, we manipulated the objective shape of the “squares” to null the perceptual pincushioning, collecting data independently at various set speeds. On each trial, the rotation rate was randomly selected from the range 0.2, 0.4, 0.5, 0.6, 0.8, and 1.0 rev/s. The observer struck designated keys to change the rotating shape along a continuum from pincushion to barrel, until satisfied that the shape looked square, neither pincushion- nor barrel-distorted. Striking the space key recorded the setting for later analysis offline and randomized the starting shape, automatically starting the next trial. Thus, observers set the physical barrel distortion to null the illusory pincushion distortion.

Corresponding author:

Sae Kaneko, Department of Psychology, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0109, USA.

Email: sakaneko@ucsd.edu



Creative Commons CC-BY: This article is distributed under the terms of the Creative Commons Attribution 3.0 License (<http://www.creativecommons.org/licenses/by/3.0/>) which permits any use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).




Figure 1. Pincushion distortion (concave sides) and barrel distortion (convex sides).




Figure 2. Pincushioning increased with rotation rates and was nulled by increasing amounts of barrel distortion. (Mean \pm 1 SE of 3 Os.)

Results are shown in Figure 2 (3 Os \times 10 settings). This plot reveals that increasing the rotation rate increased the perceptual pincushion distortion, with a 6% convexity nulling the pincushioning effect at 1.0 rev/s.

Ansbacher (1944) reported that a rotating arc looked apparently shorter along its direction of motion. This was confirmed by Stanley (1968); Anstis, Stürzel, and Spillmann (1999); and Geremek, Stürzel, da Pos, and Spillmann (2002). These authors showed that all arcs contract through equal angles at a given speed, no matter what their radii. However, whereas rotating arcs simply look shorter, our rotating squares actually appeared to change shape.

Figure 3 shows how contractions along a curved circular path can make straight lines look curved. The square in Figure 3(a) has its center at O, with its corners Q, S lying on the outer, dashed circle and the mid-side points P, R, T lying on the inner circle. The width of the square, POT, forms a straight line, thus an angle of 180°. Suppose that rapid rotation contracts the arcs of all circles, so that the radii close up like a lady's fan until the angle POT shrinks to (say) 120°. The corners Q, S and the mid-side point P, R, T are constrained to




Figure 3. Plots of (a) an undistorted square. (b) Compression distorts the square into an apparent pincushion. Note that (b) shows upper half of square only.

remain on the outer and inner circles, respectively. This will bend the previously straight side of the square QRS into a concave curve. It would be a geometric impossibility for these perceptual distortions to happen simultaneously all “around the clock.” But many illusions are inconsistent with any physical reality. For instance, the Fraser spiral illusion (Fraser, 1908) is basically a physical circle made of twisted rope. The brain uncritically accepts local votes that falsely indicate that the circle is a spiral and makes no global check that might reveal the presence of a true circle. Here, in a similar way, the brain accepts the local pincushion distortions that would be inconsistent with any physical polygon. The geometrical simulation in Figure 3 shows how tangential compression of the square makes its corners peaky and its sides concave—just like a perceptual pincushion.

Acknowledgements

The authors thank Megan Lao, Allison Lee, Jeremy Lopez, Jim Xu, and Jiajun Yuan for their assistance.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Stuart Anstis was supported by a grant from the UCSD Department of Psychology. Sae Kaneko is a research fellow of Japan Society for the Promotion of Science (supported by JSPS KAKENHI Grant number 15J03815).

Supplementary Material

The online movie 1 is available at <http://ipe.sagepub.com/supplemental>

References


Ansbacher, H. L. (1944). Distortion in the perception of real movement. *Journal of Experimental Psychology*, 34, 1–23.

- Anstis, S., Stürzel, F., & Spillmann, L. (1999). Spatial distortions in rotating radial figures. *Vision Research*, 39, 1455–1463.
- Fraser, J. (1908). A new illusion of visual direction. *British Journal of Psychology*, 2, 307–320.
- Geremek, A., Stürzel, F., da Pos, O., & Spillmann, L. (2002). Masking, persistence, and transfer in rotating arcs. *Vision Research*, 42, 2509–2519.
- Stanley, G. (1968). Apparent length of a rotating arc-line as a function of speed of rotation. *Acta Psychologica*, 28, 398–403.

Author Biographies



Stuart Anstis was born in England and was a scholar at Winchester and at Corpus Christi College, Cambridge. He took his Ph.D. at Cambridge with Prof. Richard Gregory. He has taught at the University of Bristol, UK, and at York University, Toronto, Canada. Since 1991 he has taught at the University of California, San Diego (UCSD). He has been a visiting scientist at the Smith-Kettlewell Institute, San Francisco, the San Francisco Exploratorium, and at IPRI in Japan. He has published about 170 papers on perception. He has been a Humboldt Fellow, a Visiting Fellow at Pembroke College, Oxford, and a recipient of the Kurt Koffka Medal.



Sae Kaneko received BA, MA, and PhD from The University of Tokyo. She is currently a Research Fellow of Japan Society for the Promotion of Science.